

Alcohol, Boating and Water recreation

- Alcohol is involved in 30 – 50% of all adolescent and adult drownings.^{1,2}
- Alcohol was involved in one-third of US boating fatalities in 2004. (US Boat Stats US Coast Guard)
- In Finland between 1987 - 1995, more than 63% of accidental boating and personal watercraft fatalities were associated with alcohol use.³
- Alcohol consumption by a parent or guardian may cause a lapse of supervision that contributes to child drowning.⁴
- In a New York state study of boating fatalities, boaters with a blood alcohol level of just 100 mg/dl (.01) or greater had a 16 times greater risk of drowning than those with no alcohol.⁵
- The risk of death for both operators and passengers is increased significantly with increased blood alcohol level. Therefore,

alcohol and boating prevention efforts should target not only operators but passengers as well, many of whom die when they fall intoxicated overboard when the boat is not moving.⁶

- Alcohol and drug use are proven risk factors in personal watercraft operation.⁷
- A study of teenage diving victims with severe spinal injuries also showed that alcohol was a factor in 49% of the cases.⁸
- In a study of open water diving injuries, risk factors included diving from a pier or dock, lack of familiarity with water and use of alcohol.⁹
- Experimental studies on diving while drinking show that divers with a blood alcohol count of 40 mg/dl dive significantly deeper, to a level risking spinal cord injury, although they themselves notice no difference in the depth.¹⁰

References

1. American Academy of Pediatrics. Prevention of drowning in infants, children and adolescents. Pediatrics 2003; 112:
 2. Smith GS, Brenner RA. The changing role of drowning for adolescents in the US and effective control strategies. Adolescent Medicine 1995; 6: 135- 170.
 3. Lunetta P, Penttilä A, Sarna S. Water traffic accidents, drowning and alcohol in Finland, 1969 – 1995. International Journal of Epidemiology. 1998, 27: 1038 – 1043.
 4. WHO. Facts about injuries: drowning. Department of Injuries and Violence Prevention. Geneva. Available at URL: www.who.int/violence_injury_prevention
 5. Browne ML, Lewis-Michl EL, Stark AD. Comment in: Watercraft-related drownings among New York State residents, 1988-1994. Public Health Rep. 2004 Mar-Apr;119(2):112-3; author reply 113.
 6. Smith GS, Keyl PM, Hadley JA, et al. Drinking and recreational boating fatalities: a population-based case-control study. JAMA 2001; 286: 2974 80.
 7. Van der Sman C, van Marle A, Eckhardt J, van Aken D. Risks of certain sports and recreational activities in the EU: the role of services. Consumer Safety Institute, The Netherlands, 2003; 128 p.
 8. DeVivo MJ, Sekar P. Prevention of spinal cord injuries that occur in swimming pools. Spinal Cord 1997; 35 (8): 509 – 515.
 9. Branche CM, Sniezek JE, Sattin RW, et al. Water-recreation-related spinal injuries: risk factors in natural bodies of water. Accidental Anal Prevention 1991; 23 (1): 13 – 17.
 10. Perrine MW, Mundt JC, Weiner RI. When alcohol and water don't mix: diving under the influence. Journal Study of Alcohol 1994; 55 (5): 517 – 524.
- (This fact sheet has been reviewed by EUROCARE which was formed in 1990 as an alliance of voluntary and non-governmental organisations concerned with the impact of the European Union on alcohol policy in Member States)*

The European Child Safety Alliance is a Programme of EuroSafe and is hosted by the Consumer Safety Institute in the Netherlands

PO Box 75169 1070 AD Amsterdam The Netherlands Tel. +31 20 511 45 13 Fax +31 20 511 45 10
secretariat@childsafteyeurope.org <http://www.childsafetyeurope.org>



FACTS

With the support
of the European
Commission



EUROPEAN
child
SAFETY ALLIANCE

EuroSafe